

Antiferromagnetic to valence-bond-solid transitions in two-dimensional SU(N) Heisenberg models with multi-spin interactions

Jie Lou,¹ Anders W. Sandvik,¹ and Naoki Kawashima²

¹*Department of Physics, Boston University, 590 Commonwealth Avenue, Boston, Massachusetts 02215*

²*Institute for Solid State Physics, The University of Tokyo, Kashiwa, Chiba, Japan*

(Dated: August 5, 2009)

We study two-dimensional Heisenberg antiferromagnets with additional multi-spin interactions which can drive the system into a valence-bond solid state. For standard SU(2) spins, we consider both four- and six-spin interactions. We find continuous quantum phase transitions with the same critical exponents. Extending the symmetry to SU(N), we also find continuous transitions for $N = 3$ and 4. In addition, we also study quantitatively the cross-over of the order-parameter symmetry from Z_4 deep inside the valence-bond-solid phase to U(1) as the phase transition is approached.

PACS numbers: 75.10.Jm, 75.10.Nr, 75.40.Mg, 75.40.Cx

Two-dimensional quantum spin system with non-magnetic ground states have been at the forefront of condensed matter physics for more than two decades [1, 2, 3, 4]. Frustrated system have been investigated intensely [5], but large-scale unbiased computational studies of their ground states are not possible, due to the “sign problems” hampering quantum Monte Carlo (QMC) methods [6]. It was recently realized that one prominent class of non-magnetic states—valence-bond solids (VBSs)—can be accessed also without frustration, by adding certain multi-spin interactions to the standard $S = 1/2$ Heisenberg antiferromagnet [7]. These models enable detailed QMC studies of the antiferromagnetic (AF) to VBS quantum phase transition. It has been argued that this transition is associated with spinon deconfinement (hence the term deconfined quantum criticality) and should, due to subtle quantum interference effects, be continuous [3]. This scenario violates the “Landau rule”, according to which a direct transition between states breaking unrelated symmetries should be generically first-order.

The theory of deconfined quantum criticality has generated a great deal of interest, as well as controversy [7, 8, 9, 10, 11, 12, 13, 14, 15]. Numerical studies of a Heisenberg hamiltonian with 4-spin interactions are generally in good agreement with the theory, showing a continuous transition with dynamic exponent $z = 1$, large spin correlation exponent η_s , and an emergent U(1) symmetry [7, 8, 9]. Arguments for a first-order transition have also been put forward [11, 14], based on numerical studies of lattice versions of the field-theory proposed [3] to capture the AF–VBS transition. Other, similar studies reach different conclusions, however [13]. Further studies are thus called for.

In this Letter, we advance computational studies of the AF–VBS transition in two different ways. First, we consider the $S = 1/2$ Heisenberg model including 4-spin and 6-spin interactions. The unperturbed Heisenberg model




FIG. 1: (Color online) Interactions involving p singlet projectors (illustrated by ovals enclosing two sites) on the square lattice. The two-spin ($p = 1$) interaction J is the Heisenberg exchange. Higher-order Q_p terms with $p = 2$ and 3 are considered here. All translations and 90° rotations of the site groupings shown here are included in the hamiltonian.

is defined by the hamiltonian

$$H_1 = J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j = -J \sum_{\langle ij \rangle} C_{ij} + \frac{L^2 J}{2}, \quad (1)$$

where $\langle ij \rangle$ denotes nearest neighbors on a periodic square lattice with L^2 sites and

$$C_{ij} = \frac{1}{4} - \mathbf{S}_i \cdot \mathbf{S}_j \quad (2)$$

is the two-spin singlet projector. In the “J-Q” model introduced in [7], the following term is added to H_1 ;

$$H_2 = -Q_2 \sum_{\langle ijkl \rangle} C_{kl} C_{ij}. \quad (3)$$

The spin pairs ij and kl are located on adjacent corners of a 4-site plaquette, as illustrated in Fig. 1. We denote the strength of the 4-spin term Q_2 , with the subscript indicating two singlet projectors, and also consider a similar term with three stacked singlet projectors,

$$H_3 = -Q_3 \sum_{\langle ijklmn \rangle} C_{mn} C_{kl} C_{ij}, \quad (4)$$

as also illustrated in Fig. 1. Using an improved version [16] of a ground-state QMC method operating in the valence-bond basis [17], we have studied the J- Q_2 and




FIG. 2: (Color online) Finite-size scaling of the squared AF and VBS order parameters of the J-Q₂ and J-Q₃ models.

J-Q₃ models on lattices with L up to 64. We find critical AF–VBS points with the same set of exponents for both models, providing additional evidence of a universal deconfined critical point in this class of systems.

In a second development, we have studied SU(N) symmetric versions of the J-Q₂ model, in the representation of the spin operators previously used in mean-field [2] and QMC calculations [18] of the SU(N) Heisenberg model. We find continuous AF–VBS transitions also for $N = 3$ and 4 (whereas for $N > 4$ the system is VBS ordered for all $Q_2 > 0$ [18, 19]).

An open problem in previous studies of the J-Q₂ model was that the order parameter distribution inside the VBS phase did not show the expected 4-fold symmetry. Instead, the distribution was always U(1) symmetric [7, 9]. An emergent U(1) symmetry close to criticality is indeed predicted by the field theory [3] as a consequence of a dangerously irrelevant operator, but deep inside the VBS phase the order parameter should exhibit Z₄ symmetry (which has been observed in other quantum models [19, 20]). With the J-Q₃ model and the $N > 2$ versions of the J-Q₂ model, we can now reach sufficiently deep inside the VBS phase to observe the expected U(1)–Z₄ cross-over. We present quantitative finite-size scaling results for the exponent governing the cross-over.

For all the models, we compute the square of the staggered magnetization, $M^2 = \langle \mathbf{M} \cdot \mathbf{M} \rangle$, where

$$\mathbf{M} = \frac{1}{L^2} \sum_{x,y} (-1)^{x+y} \mathbf{S}_{x,y} \quad (5)$$

is the operator of the AF (spin) order parameter. We

model, symmetry	η_s	η_d	ν	a_4
J-Q ₂ , SU(2)	0.35(2)	0.20(2)	0.67(1)	—
J-Q ₃ , SU(2)	0.33(2)	0.20(2)	0.69(2)	1.20(5)
J-Q ₂ , SU(3)	0.38(3)	0.42(3)	0.65(3)	1.6(2)
J-Q ₂ , SU(4)	0.42(5)	0.64(5)	0.70(2)	1.5(2)

TABLE I: Critical exponent for all the models studied. The cross-over exponent a_4 cannot be determined for the SU(2) J-Q₂ model, because no cross-over is observed for $L \leq 64$.

define the columnar VBS order parameter in terms of nearest-neighbor (dimer) correlators

$$D_x = \frac{1}{L^2} \sum_{x,y} (-1)^x \mathbf{S}_{x,y} \cdot \mathbf{S}_{x+1,y}, \quad (6)$$

and D_y defined analogously. We compute the square $D^2 = \langle D_x^2 + D_y^2 \rangle$ and also study the probability distribution $P(D_x, D_y)$, with D_x and D_y evaluated in the configurations generated in the QMC sampling (as in [7]). To extract the critical points and exponents, we use standard finite-size scaling forms for the order parameters,

$$M^2 = L^{1+\eta_s} F_s([q - q_c] L^{1/\nu}), \quad (7)$$

$$D^2 = L^{1+\eta_d} F_d([q - q_c] L^{1/\nu}), \quad (8)$$

where η_s and η_d are the exponents governing the spin and dimer correlation functions, respectively, at criticality (the anomalous dimensions) and $1 + \eta_{s,d} = 2\beta_{s,d}/\nu$. Here we assume a dynamic exponent $z = 1$, in accord with previous studies of the J-Q₂ model [7, 8], and use a single correlation length exponent ν , as in the theory [3].

We first present results for the SU(2) models. Defining coupling ratios $q = Q_p/(J + Q_p)$, we find critical points $q_c = 0.961(1)$ for $p = 2$ and $q_c = 0.600(5)$ for $p = 3$. The former agrees with previous estimates [7, 8, 9]. Standard data collapse plots according to (7) and (8) are shown in Fig. 2. The critical exponents are listed on the first two lines of Table I. Here it is very significant that all the exponents are the same for the two models. This supports the notion of a universal deconfined quantum critical point. Note that the order parameters decay as $L^{1+\eta_{s,d}}$ at the common critical point $q = q_c$. At a first-order transition, the order parameters should instead be size-independent at q_c , due to phase coexistence.

Comparing with previous results for the J-Q₂ model, the results for smaller systems in [7] were consistent with $\eta_s = \eta_d$ (with a value between those found here), but the present results for larger systems clearly show that the spin and dimer exponents are different. The theory does not make any specific predictions for a relationship between η_s and η_d , and they can be expected to be different. The exponents η_s and ν are in good agreement with values obtained using finite-temperature scaling in [8] (where η_d was not determined).

Next, we discuss the J-Q₂ model generalized to SU(N)




FIG. 3: (Color online) Scaling of the spin and dimer order parameters of the SU(3) and SU(4) J-Q₂ models.

spins. Considering first the Heisenberg model, the hamiltonian can be written as

$$H_{\text{SU}(N)} = \frac{J}{N} \sum_{\langle ij \rangle} \mathbf{S}_i^{\alpha\beta} \mathbf{S}_j^{\beta\alpha} = -J \sum_{\langle ij \rangle} C_{ij} + \frac{2JL^2}{N^2}, \quad (9)$$

where $\mathbf{S}_i^{\alpha\beta}$ is the generator of the SU(N) algebra, with $\alpha, \beta = 1, 2, \dots, N$ the different “colors”, and C_{ij} is the generalization of Eq. (2) to SU(N). As in [18], we focus on the simplest case, where the spins on sublattice A are expressed in the fundamental representation (i.e., with a single-box Young tableau). Spins on sublattice B are SU(N) conjugates (dual representation) of those on A (a Young tableau with one column and $N - 1$ rows). The states in this representation can be written in terms of permutations P of the boxes, with

$$|\bar{\alpha}\rangle_j \equiv \frac{1}{\sqrt{(N-1)!}} \sum_P (-1)^P |P(2)P(3) \dots P(N)\rangle_j, \quad (10)$$

with $\alpha = 1, 2, \dots, N$ and $P(1) = \alpha$. An SU(N) singlet of spins i and j on different sublattices is given by

$$|\text{singlet}\rangle_{ij} \equiv \frac{1}{\sqrt{N}} \sum_{\alpha=1}^N |\alpha\rangle_i \otimes |\bar{\alpha}\rangle_j. \quad (11)$$

QMC algorithms using these SU(N) spins in the valence-bond basis are simple generalizations of the SU(2) case [16, 17, 21]. Instead of spins $\uparrow$ and $\downarrow$ for SU(2), there are N colors, and, thus, N states of the space-time loops




FIG. 4: (Color online) Dimer order distribution $P(D_x, D_y)$ for $L = 32$ systems. The left panels are for the J-Q₃ model at $q = 0.635$ (a) and $q = 0.85$ (b), and the right panels are for the SU(3) J-Q₂ model at $q = 0.45$ (c) and $q = 0.65$ (d).

in the loop-algorithm [16]. The off-diagonal matrix elements of the singlet projection operators are $1/N$ instead of $1/2$, and the overlap of two valence-bond states is generalized to $N^{n_o - L^2/2}$, where n_o is the number of loops in the transposition graph. The 4- and 6-spin terms (3) and (4) are written explicitly using products of singlet projectors and have obvious generalizations to SU(N).

Our results for the SU(3) and SU(4) versions of the J-Q₂ model are consistent with continuous AF-VBS critical points, with no signs of first-order behavior. The critical points are $q_c = 0.335(2)$ and $q_c = 0.082(2)$ for $N = 3$ and 4, respectively. Scaling plots giving the critical exponents are shown in Fig. 3 and numerical values are listed in Table I. It can be noted that, as a function of N , ν does not change appreciably, η_s increase slowly, and η_d increases significantly. An increasing spin exponent is consistent with $\eta_s = 1$ in the $N \rightarrow \infty$ theory [3]. As already noted, there are no predictions for η_d . We could, in principle, consider still higher N , but with $J > 0$ the system is always in the VBS state for $N = 5$ and higher [18, 19]. A transition could presumably be reached for $J < 0$, but this causes QMC sign problems.

The dimer order distribution $P(D_x, D_y)$ can be used to investigate the VBS order parameter symmetry [7, 18]. As shown in Fig. 4, for large q the robust VBSs in the SU(2) J-Q₃ model and the SU(3) and SU(4) versions of the J-Q₂ model result in histograms with clearly visible columnar Z_4 features (i.e., peaks on the D_x and D_y axis, as opposed to 45° degree rotated histograms expected for a plaquette state). However, in the SU(2) J-Q₂ model the histograms are ring-shaped for all system sizes currently accessible, even in the extreme case of $q = 1$ ($J = 0$). In all cases, we see U(1) symmetric histograms as the critical point is approached, in agreement with one of the salient features of deconfined quantum criticality [3].




FIG. 5: (Color online) Finite-size scaling of the square of the anisotropic order parameter D_4 in the J- Q_3 model (upper panel) and SU(3) J- Q_2 model (lower panel).

Defining an order parameter sensitive to the symmetry,

$$\begin{aligned} D_4^2 &= \int dD_x dD_y P(D_x, D_y) (D_x^2 + D_y^2) \cos(4\theta) \\ &= \int dr \int_0^{2\pi} d\theta P(r, \theta) r^3 \cos(4\theta), \end{aligned} \quad (12)$$

where θ is the angle corresponding to a point (D_x, D_y) , we proceed as in [22] (which deals with a classical system with a dangerously irrelevant perturbation) to extract the exponent governing the length-scale Λ of the Z_4 -U(1) cross-over (and the spinon confinement). Z_4 features should appear for $L > \Lambda$, which is predicated [3] to scale as $\Lambda \sim \xi^{a_4}$ where ξ is the correlation length and $a_4 > 1$. We analyze D_4 assuming the scaling form [22];

$$D_4^2 = L^{1+\eta_d} F_4(qL^{1/a_4\nu}). \quad (13)$$

This form describes the cross-over, as shown in Fig. 5 in two cases. The values of a_4 are listed in Table I. The large error bars reflect slow evolution of the VBS angle in the QMC simulations. It is nevertheless clear that $a_4 > 1$ (and increasing with N), reflecting emergent U(1) symmetry due to a dangerously irrelevant perturbation.

The results presented here support deconfined quantum criticality. Although one can still, in principle, not completely rule out very weakly first-order transitions based on these calculations, the universal behavior for the two SU(2) models makes this less likely. The common exponents for the J- Q_2 and J- Q_3 models at the very least suggest close proximity to a universal critical point.

The detailed information now available from QMC simulations should be useful to further advance the theory.

In a very interesting experimental development, Ito *et al.* have recently measured the spin-lattice relaxation rate $1/T_1$ in a layered organic compound which seems to be near-critical [23]. It has been argued that, in spite of the triangular lattice, the AF-VBS transition in this kind of system should be in the same class of deconfined quantum critical points discussed here [24]. The exponent η_s governs the temperature scaling of $1/T_1$, and the value $\eta_s \approx 0.35$ is in excellent agreement with the experiment over a wide range of temperatures. Further experiments should elucidate the nature of the ground state and whether it is indeed close to a deconfined quantum-critical point.

Acknowledgements—We would like to thank R. Kaul and A. Vishwanath for useful discussions. This work was supported by NSF grant No. DMR-0803510 (J. L. and A. W. S) and MEXT Grant-in-Aid for Scientific Research [(B)(19340109) and Priority Area “Novel States of Matter Induced by Frustration” (19052004)] (N. K.).

-
- [1] P. W. Anderson, *Science* **235** 1196 (1987).
 - [2] N. Read and S. Sachdev, *Phys. Rev. Lett.* **62**, 1694 (1989).
 - [3] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, *Science* **303**, 1490 (2004); M. Levin and T. Senthil, *Phys. Rev. B* **70**, 220403 (2004).
 - [4] S. Sachdev, *Nature Physics* **4**, 173 (2008).
 - [5] H. T. Diep (editor), *Frustrated Spin Systems* (World Scientific, 2005).
 - [6] P. Henelius and A. W. Sandvik, *Phys. Rev. B* **62**, 1102 (2000); F.-J. Jiang, F. Kämpfer, M. Nyfeler, and U.-J. Wiese, *Phys. Rev. B* **78**, 214406 (2008).
 - [7] A. W. Sandvik, *Phys. Rev. Lett.* **98**, 227202 (2007).
 - [8] R. G. Melko and R. K. Kaul, *Phys. Rev. Lett.* **100**, 017203 (2008); R. K. Kaul and R. G. Melko, *Phys. Rev. B* **78**, 014417 (2008).
 - [9] F.-J. Jiang, M. Nyfeler, S. Chandrasekharan, and U.-J. Wiese, *J. Stat. Mech.*, P02009 (2008). Here a first-order transition in the J-Q model is argued, but large statistical errors make the data open to other interpretations.
 - [10] Z. Weihong, O. P. Sushkov, and J. Oitmaa, *Phys. Rev. B* **73**, 184420 (2006).
 - [11] A.B. Kuklov, N.V. Prokof'ev, B.V. Svistunov and M. Troyer, *Annals of Physics* **321**, 1602 (2006).
 - [12] F. S. Nogueira, S. Kragset, and A. Sudbø, *Phys. Rev. B* **76**, 220403 (2007).
 - [13] O. I. Motrunich and A. Vishwanath, arXiv:0805.1494.
 - [14] A. Kuklov *et al.*, *Phys. Rev. Lett.* **101**, 050405 (2008).
 - [15] F. S. Nogueira, *Phys. Rev. D* **79**, 105007 (2009).
 - [16] A. W. Sandvik and H. G. Evertz, arXiv:0807.0682.
 - [17] A. W. Sandvik, *Phys. Rev. Lett.* **95**, 207203 (2005).
 - [18] N. Kawashima and Y. Tanabe, *Phys. Rev. Lett.* **98**, 057202 (2007).
 - [19] K. S. D. Beach, F. Alet, M. Mambrini, and S. Capponi, arXiv:0812.3657.
 - [20] J. Lou and A. W. Sandvik, arXiv:0811.0837.

- [21] An equivalent way to generalize to $N > 2$ is to consider spin $S = (N - 1)/2$ models including higher-order exchange operators rendering the hamiltonian $SU(N)$ symmetric. This approach was developed for the Heisenberg model in [19].
- [22] J. Lou, A. W. Sandvik, and L. Balents, Phys. Rev. Lett. **99**, 207203 (2007).
- [23] T. Itou et al., Phys. Rev. B **77**, 104413 (2008).
- [24] C. Xu and S. Sachdev, Phys. Rev. B **79**, 064405 (2009).